arXiv:1402.4373v1 [math.CO] 18 Feb 2014

A COMMENT ON: "FURTHER RESTRICTIONS ON THE STRUCTURE OF FINITE DCI-GROUPS"

EDWARD DOBSON, JOY MORRIS, AND PABLO SPIGA

Abstract. A finite group R is a DCI-group if, whenever S and T are subsets of R with the Cayley graphs $\mathrm{Cay}(R,S)$ and $\mathrm{Cay}(R,T)$ isomorphic, there exists an automorphism φ of R with $S^\varphi = T$.

The classification of DCI-groups is an open problem in the theory of Cayley graphs and is closely related to the isomorphism problem for graphs. This paper is a contribution towards this classification, as we show that every dihedral group of order $6p$, with $p \geq 5$ prime, is a DCI-group. This corrects and completes the proof of [5, Theorem 1.1] as observed by the reviewer [3].

1. Introduction

Let R be a finite group and let S be a subset of R. The *Cayley digraph* of R with connection set S, denoted $\mathrm{Cay}(R,S)$, is the digraph with vertex set R and with (x,y) being an arc if and only if $xy^{-1} \in S$. Now, $\mathrm{Cay}(R,S)$ is said to be a *Cayley isomorphic* digraph, or DCI-*graph* for short, if whenever $\mathrm{Cay}(R,S)$ is isomorphic to $\mathrm{Cay}(R,T)$, there exists an automorphism φ of R with $S^\varphi = T$. Clearly, $\mathrm{Cay}(R,S) \cong \mathrm{Cay}(R,S^\varphi)$ for every $\varphi \in \mathrm{Aut}(R)$ and hence, loosely speaking, for a DCI-graph $\mathrm{Cay}(R,S)$ deciding when a Cayley digraph over R is isomorphic to $\mathrm{Cay}(R,S)$ is theoretically and algorithmically elementary; that is, the solving set for $\mathrm{Cay}(R,S)$ is reduced to simply $\mathrm{Aut}(R)$ (for the definition of solving set see for example [6, 7]). The group R is a DCI-*group* if $\mathrm{Cay}(R,S)$ is a DCI-graph for every subset S of R. Moreover, R is a CI-group if $\mathrm{Cay}(R,S)$ is a DCI-graph for every inverse-closed subset S of R. Thus every DCI-group is a CI-group.

Throughout this paper, p will always denote a prime number.

In order to obtain new and severe constrains on the structure of a DCI-group, the authors of [5] considered the problem of determining which Frobenius groups R of order $6p$ are DCI-groups. They were in fact interested in the more specific case of Frobenius groups of order $6p$ with Frobenius kernel of order p; this is clear from their analysis and their proofs, but is not specified in the statement of [5, Theorem 1.1]. The proof of their theorem as stated is therefore incomplete, as observed by Conder [3]. The aim of this paper is to fix this discrepancy by completing the analysis of which Frobenius groups of order $6p$ are DCI-groups, hence completing the proof of [5, Theorem 1.1] as the authors stated it.

2010 *Mathematics Subject Classification.* 20B10, 20B25, 05E18.

Key words and phrases. Cayley graph, isomorphism problem, CI-group, dihedral group.

Address correspondence to P. Spiga, E-mail: pablo.spiga@unimib.it

The second author is supported in part by the National Science and Engineering Research Council of Canada.

An elementary computation yields that if R is a Frobenius group of order $6p$ with Frobenius kernel whose order is not p, then R is isomorphic to the alternating group on four symbols Alt(4) (and $p = 2$), or to the quasidihedral group $\langle (1,2,3),(4,5,6),(2,3)(5,6)\rangle$ (and $p = 3$), or to the dihedral group of order $6p$. A routine computer-assisted computation shows that Alt(4) is a DCI-group and $\langle (1,2,3),(4,5,6),(2,3)(5,6)\rangle$ is not a DCI-group. Moreover, as is observed in [3], $\langle (1,2,3),(4,5,6),(2,3)(5,6)\rangle$ is a CI-group. Therefore in order to complete the analysis of Frobenius groups of order $6p$, we only need to consider dihedral groups of order $6p$.

Theorem 1.1. *Let p be a prime number and let R be the dihedral group of order $6p$. Then R is a* DCI*-group if and only if $p \geq 5$, and R is a* CI*-group if and only if $p \geq 3$.*

The structure of the paper is straightforward. In Section 2, we consider the case $p \leq 5$. In Section 3, we provide some preliminary definitions and our main tool. In Section 4 we introduce some notation and we divide the proof of Theorem 1.1 into four cases, which we then study in turn in Sections 5–8.

2. Small groups: $p \leq 5$

Lemma 2.1. *Let p be a prime with $p \leq 5$ and let R be the dihedral group of order $6p$. Then R is a* DCI*-group if and only if $p = 5$, and R is a* CI*-group if and only if $p \neq 2$.*

Proof. The proof follows from a computer computation with the invaluable help of the algebra system `magma` [2]. Let $R_p = \langle a, b \mid a^{3p} = b^2 = (ab)^2 = 1\rangle$ be the dihedral group of order $6p$. Here we simply prove that R_2 is not a CI-group and that R_3 is not a DCI-group.

For $p = 2$, the graphs $\mathrm{Cay}(R_2, \{b, a^3\})$ and $\mathrm{Cay}(R_2, \{b, a^3b\})$ are both isomorphic to the disjoint union of three cycles of length 4. As a^3 is the only central involution of R_2, there exists no automorphism of R_2 mapping $\{b, a^3\}$ to $\{b, a^3b\}$.

For $p = 3$, the digraphs $\mathrm{Cay}(R_3, \{a, a^4, a^6, a^7\})$ and $\mathrm{Cay}(R_3, \{a^2, a^5, a^6, a^8\})$ are isomorphic and a computation shows that there exists no automorphism of R_3 mapping $\{a, a^4, a^6, a^7\}$ to $\{a^2, a^5, a^6, a^8\}$. □

Given that the (di)graphs we described in this proof are not connected, it is worth observing that a group R is a CI-group if and only if every pair of connected isomorphic Cayley graphs on R are isomorphic via an automorphism of R. This is because the complement of a disconnected graph is always connected, and the property of being a CI-graph is preserved under taking complements. A similar observation also applies to DCI-groups.

In view of Lemma 2.1 for the rest of this paper we may assume that $p \geq 7$.

3. Some basic results

Babai [1] has proved a very useful criterion for determining when a finite group R is a DCI-group and, more generally, when $\mathrm{Cay}(R, S)$ is a DCI-graph.

Lemma 3.1. *Let R be a finite group and let S be a subset of R. Then* $\mathrm{Cay}(R, S)$ *is a* DCI*-graph if and only if* $\mathrm{Aut}(\mathrm{Cay}(R, S))$ *contains a unique conjugacy class of regular subgroups isomorphic to R.*

Let Ω be a finite set and let G be a permutation group on Ω. The 2-*closure* of G, denoted $G^{(2)}$, is the set

$$\{\pi \in \mathrm{Sym}(\Omega) \mid \forall(\omega,\omega') \in \Omega^2, \text{there exists } g_{\omega\omega'} \in G \text{ with } (\omega,\omega')^\pi = (\omega,\omega')^{g_{\omega\omega'}}\},$$

where $\mathrm{Sym}(\Omega)$ is the symmetric group on Ω. Observe that in the definition of $G^{(2)}$, the element $g_{\omega\omega'}$ of G may depend upon the ordered pair (ω,ω'). The group G is said to be 2-*closed* if $G = G^{(2)}$.

It is easy to verify that $G^{(2)}$ is a subgroup of $\mathrm{Sym}(\Omega)$ containing G and, in fact, $G^{(2)}$ is the smallest (with respect to inclusion) subgroup of $\mathrm{Sym}(\Omega)$ preserving every orbital digraph of G. It follows that the automorphism group of a graph is 2-closed. Therefore Lemma 3.1 immediately yields:

Lemma 3.2. *Let R be a finite group and let R_r be the right regular representation of R in Sym(R). If, for every $\pi \in$ Sym(R), the groups R_r and R_r^π are conjugate in $\langle R_r, R_r^\pi\rangle^{(2)}$, then R is a* DCI*-group.*

Proof. Let S be a subset of R, and set $\Gamma := \mathrm{Cay}(R,S)$ and $A := \mathrm{Aut}(\Gamma)$. Observe that $R_r \leq A$ and that A is 2-closed. Let T be a regular subgroup of A isomorphic to R. Since $\langle R_r, T\rangle \leq A$, we get $\langle R_r, T\rangle^{(2)} \leq A^{(2)} = A$.

Every regular subgroup of $\mathrm{Sym}(R)$ isomorphic to R is conjugate to R_r and hence $T = R_r^\pi$, for some $\pi \in \mathrm{Sym}(R)$. By hypothesis, R_r and T are conjugate in $\langle R_r, T\rangle^{(2)}$ and so are conjugate in A. In particular, A contains a unique conjugacy class of regular subgroups isomorphic to R and Lemma 3.1 gives that R is a DCI-group. □

We will use this formulation of Babai's criterion without comment in our proof of Theorem 1.1.

4. Notation and preliminary reductions

Multiplication of permutations is on the right, so $\sigma\tau$ is calculated by first applying σ, and then τ. For the rest of this paper we let R be the dihedral group of order $6p$ and we let $\Omega := \{1,\ldots,6p\}$. Using Lemma 2.1, we may assume that $p \geq 7$ in the proof of Theorem 1.1. In what follows, we identify R with a regular subgroup of $\mathrm{Sym}(\Omega)$ isomorphic to R, that is, R acts regularly on Ω. Let $\pi \in \mathrm{Sym}(\Omega)$ and set $G := \langle R, R^\pi\rangle$. In view of Lemma 3.2, Theorem 1.1 will follow by proving that R is conjugate to R^π via an element of $G^{(2)}$.

Let R_p denote the Sylow p-subgroup of R, let P be a Sylow p-subgroup of G with $R_p \leq P$ and let T be a Sylow p-subgroup of $\mathrm{Sym}(\Omega)$ with $P \leq T$. From Sylow's theorems, replacing R^π by a suitable G-conjugate, we may assume that $R_p^\pi \leq P$. Observe that, as $p \geq 7$, the group T is elementary abelian of order p^6. Since R_p and R_p^π are acting semiregularly, their orbits on Ω must be equal to the orbits of T.

Since R_p is the unique Sylow p-subgroup of R, we see that R admits a unique system of imprimitivity $\mathcal{C}$ with blocks of size p, namely $\mathcal{C}$ consists of the R_p-orbits on Ω. Similarly, R^π admits a unique system of imprimitivity with blocks of size p, namely $\mathcal{C}^\pi$, and the system of imprimitivity $\mathcal{C}^\pi$ consists of the R_p^π-orbits on Ω. Since each of these is equal to the orbits of T on Ω, we have $\mathcal{C} = \mathcal{C}^\pi$, and $\mathcal{C}$ is R- and R^π-invariant. As $G = \langle R, R^\pi\rangle$, we get that $\mathcal{C}$ is also G-invariant. Therefore, G is conjugate to a subgroup of $\mathrm{Sym}(p) \operatorname{wr} \mathrm{Sym}(6)$. Similarly, since $\mathcal{C}$ is π-invariant, π is conjugate to an element in $\mathrm{Sym}(p) \operatorname{wr} \mathrm{Sym}(6)$.

We can use this structure to decompose the set Ω as $\Delta \times \Lambda$ with $|\Delta| = p$ and $|\Lambda| = 6$. We identify Ω with $\Delta \times \Lambda$, Δ with $\{1, \ldots, p\}$ and Λ with $\{1, \ldots, 6\}$. Write $W := \mathrm{Sym}(\Delta) \operatorname{wr} \mathrm{Sym}(\Lambda)$ and $B := \mathrm{Sym}(\Delta)^6$ the base group of W. Then for $\sigma \in \mathrm{Sym}(\Lambda)$, $(y_1, \ldots, y_6) \in B$, and $(\delta, \lambda) \in \Delta \times \Lambda$, we have

$$(\delta, \lambda)^\sigma = (\delta, \lambda^\sigma) \text{ and } (\delta, \lambda)^{(y_1,\ldots,y_6)} = (\delta^{y_\lambda}, \lambda),$$

and $W = \{\sigma(y_1, \ldots, y_6) \mid \sigma \in \mathrm{Sym}(\Lambda), (y_1, \ldots, y_6) \in B\}$. Observe that under this identification the system of imprimitivity $\mathcal{C}$ is $\{\Delta_1, \ldots, \Delta_6\}$ where $\Delta_\lambda = \Delta \times \{\lambda\}$ for every $\lambda \in \Lambda$.

Let K be the kernel of the action of G on $\mathcal{C}$, that is, $K = B \cap G$. Clearly, RK/K and $R^\pi K/K$ are regular subgroups of $\mathrm{Sym}(\Lambda)$ isomorphic to $\mathrm{Sym}(3)$. A direct inspection in $\mathrm{Sym}(\Lambda)$ shows that if A and B are regular subgroups of $\mathrm{Sym}(\Lambda)$ isomorphic to $\mathrm{Sym}(3)$, then either B is conjugate to A via an element of $\langle A, B\rangle$, or $\langle A, B\rangle = A \times B$. Summing up and applying this observation to G/K, we obtain the following reduction.

Reduction 4.1. We have

$$G \leq W \quad \text{and} \quad \pi \in W,$$

and (replacing G by a suitable W-conjugate) either

$$\frac{G}{K} = \frac{RK}{K} = \frac{R^\pi K}{K} = \langle (1,2,3)(4,5,6), (1,4)(2,6)(3,5)\rangle, \tag{1}$$

or

$$\begin{aligned} \frac{G}{K} &= \frac{RK}{K} \times \frac{R^\pi K}{K}, \\ RK/K &= \langle (1,2,3)(4,5,6), (1,4)(2,6)(3,5), \\ R^\pi K/K &= \langle (1,2,3)(4,6,5), (1,4)(2,5)(3,6)\rangle. \end{aligned} \tag{2}$$

A moment's thought gives that in case (1) we may assume that $\pi \in B$ and in case (2) we may assume that $\pi = (5,6)y$ with $y \in B$. Write $\pi := \sigma(y_1, \ldots, y_6)$ with $\sigma = 1$ or $\sigma = (5,6)$ depending on whether case (1) or (2) is satisfied. Set $y := (y_1, \ldots, y_6)$.

Let c be the cycle $(1, 2, \ldots, p)$ of length p of $\mathrm{Sym}(\Delta)$. Set

$$r_1 := (c, c, c, c, c, c),\ r_2 := (1,2,3)(4,5,6) \text{ and } r_3 := (1,4)(2,6)(3,5).$$

Replacing G by a suitable W-conjugate, we may assume that

$$R_p = \langle r_1\rangle \text{ and } R = \langle r_1, r_2, r_3\rangle. \tag{3}$$

Clearly, $\mathbf{N}_{\mathrm{Sym}(\Delta)}(\langle c\rangle) \cong \mathrm{AGL}_1(p)$ and hence $\mathbf{N}_{\mathrm{Sym}(\Delta)}(\langle c\rangle) = \langle c, \alpha\rangle = \langle c\rangle \rtimes \langle \alpha\rangle$, where α is a permutation fixing 1 and acting by conjugation on $\langle c\rangle$ as an automorphism of order $p - 1$.

As $R_p \leq T$, we see that T is generated by $c_1, c_2, \ldots, c_6$ where

$$c_1 := (c, 1, 1, 1, 1, 1), c_2 := (1, c, 1, 1, 1, 1), \ldots, c_6 := (1, 1, 1, 1, 1, c).$$

Since $R_p^\pi \leq T$ and since R_p^π is semiregular, we obtain

$$R_p^\pi = \langle (c^{\ell_1}, c^{\ell_2}, c^{\ell_3}, c^{\ell_4}, c^{\ell_5}, c^{\ell_6})\rangle,$$

with $\ell_1 = 1$ and for some $\ell_2, \ldots, \ell_6 \in \{1, \ldots, p-1\}$.

Now $r_1^\pi = (c^{y_1}, c^{y_2}, c^{y_3}, c^{y_4}, c^{y_5}, c^{y_6}) \in R_p^\pi$ and hence there exists $\ell \in \{1, \dots, p-1\}$ with $c^{y_\lambda} = c^{\ell_\lambda \ell}$, for every $\lambda \in \Lambda$. Thus $y_\lambda \in \mathbf{N}_{\mathrm{Sym}(\Delta)}(\langle c \rangle) = \langle c, \alpha \rangle$ and $y_\lambda = c^{u_\lambda}\alpha^{v_\lambda}$ for some $u_\lambda \in \{0, \dots, p-1\}$ and $v_\lambda \in \{0, \dots, p-2\}$. It follows that

$$\begin{aligned} \pi &= \sigma(c^{u_1}\alpha^{v_1}, c^{u_2}v^{\alpha_2}, \dots, c^{u_6}\alpha^{v_6}) \in \langle c, \alpha \rangle \operatorname{wr} \mathrm{Sym}(\Lambda), \\ G &\leq \langle c, \alpha \rangle \operatorname{wr} \mathrm{Sym}(\Lambda). \end{aligned} \tag{4}$$

Now $r_1 \in R \leq G$, and hence replacing π by $r_1^{-u_1}\pi$, we may assume that $u_1 = 0$. Furthermore, $(\alpha, \alpha, \alpha, \alpha, \alpha, \alpha) \in \mathbf{N}_{\mathrm{Sym}(\Omega)}(R)$, and hence replacing π by $(\alpha, \dots, \alpha)^{-v_1}\pi$, we may assume that $v_1 = 0$.

As $\langle c, \alpha \rangle \operatorname{wr} \mathrm{Sym}(\Lambda)$ has a normal Sylow p-subgroup, we get $P \trianglelefteq G$ and K/P is isomorphic to a subgroup of $\langle \alpha \rangle \times \langle \alpha \rangle \times \langle \alpha \rangle \times \langle \alpha \rangle \times \langle \alpha \rangle \times \langle \alpha \rangle$.

Next we define an equivalence relation $\equiv$ on Ω. We say that $\omega \equiv \omega'$ if $P_\omega = P_{\omega'}$. Since $P \trianglelefteq G$, we see that $\equiv$ is G-invariant. Moreover, since P is abelian, we get that P acts regularly on each of its orbits and hence $\omega \equiv \omega'$ for every ω and ω' in the same P-orbit. This shows that $\equiv$ defines a system of imprimitivity $\mathcal{E}$ for G coarser than $\mathcal{C}$. In particular, $\equiv$ consists of either 1, 2, 3 or 6 equivalence classes.

There is an equivalent definition of $\equiv$. Given $\omega \in \Delta_\lambda$ and $\omega' \in \Delta_{\lambda'}$, we have $\omega \equiv \omega'$ whenever, for every $\rho \in P$, $\rho|_{\Delta_\lambda} = 1$ if and only if $\rho|_{\Delta_{\lambda'}} = 1$ (or equivalently, $\rho|_{\Delta_\lambda}$ is a p-cycle if and only if $\rho|_{\Delta_{\lambda'}}$ is a p-cycle).

We will use the following lemma repeatedly.

Lemma 4.2. *For every $\rho \in K$ and for every $E \in \mathcal{E}$, the permutation $\rho_E : \Omega \to \Omega$, fixing $\Omega \setminus E$ pointwise and acting on E as ρ does, lies in $G^{(2)}$.*

Proof. This is Lemma 2 in [4]. (We remark that [4, Lemma 2] is only stated for graphs, but the result holds for each orbital digraph of G, and hence for $G^{(2)}$.) □

With all of this notation at our disposal we are ready to prove Theorem 1.1 with a case analysis depending on the number of $\equiv$-equivalence classes.

5. Case I: $\equiv$ has only one equivalence class

Here, $P_\omega = P_{\omega'}$ for every $\omega, \omega' \in \Omega$, hence P acts semiregularly on Ω and $|P| = p$. It follows that $P = R_p = R_p^\pi$. In particular, $\ell_1 = \cdots = \ell_6 = 1$ and $v_1 = \cdots = v_6 = 0$. Therefore $\pi = \sigma(c^{u_1}, c^{u_2}, c^{u_3}, c^{u_4}, c^{u_5}, c^{u_6})$ with $u_1 = 0$.

Suppose that $\sigma = 1$. Since $r_2, r_2^\pi \in G$, we have

$$r_2^{-1}(r_2)^\pi = (c^{-u_3+u_1}, c^{-u_1+u_2}, c^{-u_2+u_3}, c^{-u_6+u_4}, c^{-u_4+u_5}, c^{-u_5+u_6}) \in P$$

and hence $-u_3+u_1 = -u_1+u_2 = -u_2+u_3 = -u_6+u_4 = -u_4+u_5 = -u_5+u_6$. This gives $u_1 = u_2 = u_3 = 0$ and $u_4 = u_5 = u_6$. Write $u := u_4$. A similar computation gives

$$r_3^{-1}(r_3)^\pi = (c^{-u}, c^{-u}, c^{-u}, c^u, c^u, c^u) \in P.$$

Thus $u = -u$ and hence $u = 0$. Therefore $\pi = 1$ and $R^\pi = R$. It follows that R is conjugate to R^π via the identity element of $G^{(2)}$.

Suppose that $\sigma = (5, 6)$. Since $r_2, r_2^\pi \in G$, we have

$$r_2^{-1}(r_2)^\pi = (4, 5, 6)(c^{-u_3+u_1}, c^{-u_1+u_2}, c^{-u_2+u_3}, c^{-u_5+u_4}, c^{-u_6+u_5}, c^{-u_4+u_6}) \in G$$

and by taking the 3rd power we get $(c^{3(-u_3+u_1)}, c^{3(-u_1+u_2)}, c^{3(-u_2+u_3)}, 1, 1, 1) \in P$. Thus $3(-u_3 + u_1) = 3(-u_1 + u_2) = 3(-u_2 + u_3) = 0$ and since $u_1 = 0$, we have

$u_1 = u_2 = u_3 = 0$. Moreover

$$r_2(r_2)^\pi = (1,3,2)(1,1,1,c^{-u_5+u_4},c^{-u_6+u_5},c^{-u_4+u_6}) \in G$$

and by taking the 3rd power we get $(1,1,1,c^{3(-u_5+u_4)},c^{3(-u_6+u_5)},c^{3(-u_4+u_6)}) \in P$. Thus $3(-u_5+u_4) = 3(-u_6+u_5) = 3(-u_4+u_6) = 0$ and hence $u_4 = u_5 = u_6$. Write $u := u_4$. Now

$$r_3^{-1}(r_3)^\pi = (2,3)(5,6)(c^{-u},c^{-u},c^{-u},c^{u},c^{u},c^{u}) \in G$$

and by taking the 2nd power we get $(c^{-2u},c^{-2u},c^{-2u},c^{2u},c^{2u},c^{2u}) \in P$. Thus $2u = -2u$, and hence $u = 0$. It follows that $\pi = \sigma = (5,6)$ and

$$G = \langle R, R^\pi \rangle = \langle r_1, (1,2,3)(4,5,6), (1,4)(2,6)(3,5), (1,2,3)(4,6,5), (1,4)(2,5)(3,6) \rangle.$$

We claim that $\pi \in G^{(2)}$, from which the proof of this case follows. First observe that $(1,2,3)(4,5,6)(1,2,3)(4,6,5) = (1,3,2) \in G$. Also $r_3^{-1}r_3^\pi = (2,3)(5,6) \in G$, and hence (conjugating by the elements of $\langle (1,3,2) \rangle$), we see that $(1,2)(5,6)$ and $(1,3)(5,6)$ belong to G. Next, let $\omega = (\delta,\lambda)$ and $\omega' = (\delta',\lambda')$ be in Ω. If $\lambda,\lambda' \notin \{5,6\}$, then $(\omega,\omega')^\pi = (\omega,\omega')^{g_{\omega\omega'}}$ with $g_{\omega\omega'} = 1$. If $\lambda,\lambda' \in \{5,6\}$, then $(\omega,\omega')^\pi = (\omega,\omega')^{g_{\omega\omega'}}$ with $g_{\omega\omega'} = (1,2)(5,6)$. Finally, suppose that only one of λ,λ' lies in $\{5,6\}$. Let λ'' be the element of $\{\lambda,\lambda'\} \cap \{1,2,3,4\}$ and let $g_{\omega\omega'}$ be in $\{(1,2)(5,6),(1,3)(5,6),(2,3)(5,6)\}$ fixing the block $\Delta_{\lambda''}$ pointwise. Then $(\omega,\omega')^\pi = (\omega,\omega')^{g_{\omega\omega'}}$.

6. Case II: $\equiv$ has six equivalence classes

Since $\equiv$ has six equivalence classes, for every two distinct $\lambda,\lambda' \in \Lambda$, there exists an element $q \in P$ with q fixing Δ_λ pointwise and acting as the cycle c on $\Delta_{\lambda'}$. From this it follows that $P^{(2)} = T$. Next, from $T \leq G^{(2)}$, it follows that if $\gamma : \Omega \to \Omega$ is a permutation with the property that for each $\lambda \in \Lambda$, we have

- $\Delta_\lambda^\gamma = \Delta_\lambda$ and
- $\gamma|_{\Delta_\lambda} = g_\lambda|_{\Delta_\lambda}$ for some $g_\lambda \in G$ fixing Δ_λ setwise,

then $\gamma \in G^{(2)}$.

As $T = P^{(2)} \leq G^{(2)}$, replacing π by $g^{-1}\pi$ for a suitable $g \in T$, we may assume that $u_1 = u_2 = \cdots = u_6 = 0$.

For $2 \leq \lambda \leq 6$, let g_λ be the element of R that maps $(1,1)$ to $(1,\lambda)$ (so $g_2 = r_2$, etc.). Define $\gamma : \Omega \to \Omega$ by $\gamma|_{\Delta_1} = \mathrm{id}|_{\Delta_1}$, and for $2 \leq \lambda \leq 6$,

$$\gamma|_{\Delta_\lambda} = \left((g_{\lambda^\sigma}^\pi)^{-1} g_\lambda\right)|_{\Delta_\lambda}.$$

By the observations we made in the first paragraph of this case, $\gamma \in G^{(2)}$. Careful computations show that $(r_1^\pi)^\gamma = r_1$. Thus, $(R_p^\pi)^\gamma = R_p$. We now see that after conjugating R^π by γ we are in Case I and can complete the proof as before.

7. Case III: $\equiv$ has two equivalence classes

The $\equiv$-equivalence classes are blocks of imprimitivity for G of size $3p$ and are a union of P-orbits. The only system of imprimitivity for G/K with blocks of size 3 is $\{\{1,2,3\},\{4,5,6\}\}$. Therefore the two $\equiv$-equivalence classes are $\Delta_1 \cup \Delta_2 \cup \Delta_3$ and $\Delta_4 \cup \Delta_5 \cup \Delta_6$. By Lemma 4.2 applied to $\rho = r_1$, $(c,c,c,1,1,1),(1,1,1,c,c,c) \in G^{(2)}$.

Replacing π by $g^{-1}\pi$ for a suitable $g \in G^{(2)}$, we may assume that $u_4 = 0$. As $R_p^\pi \leq P$, we get $\ell_1 = \ell_2 = \ell_3$ and $\ell_4 = \ell_5 = \ell_6$. It follows that $v_1 = v_2 = v_3 = 0$ and $v_4 = v_5 = v_6$. Write $\beta := \alpha^{v_4}$. Therefore $\pi = \sigma(1,c^{u_2},c^{u_3},\beta,c^{u_5}\beta,c^{u_6}\beta)$.

Suppose that $\sigma = 1$. We have

$$r_2^{-1}(r_2)^\pi = (c^{-u_3}, c^{u_2}, c^{-u_2+u_3}, \beta^{-1}c^{-u_6}\beta, \beta^{-1}c^{u_5}\beta, \beta^{-1}c^{-u_5+u_6}\beta) \in P$$

and hence $-u_3 = u_2 = -u_2 + u_3$ and $-u_6 = u_5 = -u_5 + u_6$. This gives $u_2 = u_3 = 0$ and $u_5 = u_6 = 0$, that is, $\pi = (1, 1, 1, \beta, \beta, \beta)$. A similar computation gives

$$r_3^{-1}(r_3)^\pi = (\beta^{-1}, \beta^{-1}, \beta^{-1}, \beta, \beta, \beta) \in K.$$

Applying Lemma 4.2 with $E := \Delta_4 \cup \Delta_5 \cup \Delta_6$ and $\rho := r_3^{-1}(r_3)^\pi$, we get $(1, 1, 1, \beta, \beta, \beta) \in G^{(2)}$, that is, $\pi \in G^{(2)}$, from which the proof follows.

Suppose that $\sigma = (5, 6)$. Since $r_2, r_2^\pi \in G$, we have

$$r_2^{-1}(r_2)^\pi = (4, 5, 6)(c^{-u_3}, c^{u_2}, c^{-u_2+u_3}, \beta^{-1}c^{-u_5}\beta, \beta^{-1}c^{-u_6+u_5}\beta, \beta^{-1}c^{u_6}\beta) \in G$$

and by taking the 3^{rd} power we get $(c^{-3u_3}, c^{3u_2}, c^{3(-u_2+u_3)}, 1, 1, 1) \in P$. Thus $-3u_3 = 3u_2 = 3(-u_2 + u_3)$ and hence $u_1 = u_2 = u_3 = 0$. Moreover

$$r_2(r_2)^\pi = (1, 3, 2)(1, 1, 1, \beta^{-1}c^{-u_5}\beta, \beta^{-1}c^{-u_6+u_5}\beta, \beta^{-1}c^{u_6}\beta) \in G$$

and by taking the 3^{rd} power we get $(1, 1, 1, \beta^{-1}c^{-3u_5}\beta, \beta^{-1}c^{3(-u_6+u_5)}\beta, \beta^{-1}c^{3u_6}\beta) \in P$. Thus $-3u_5 = 3(-u_6 + u_5) = 3u_6$ and hence $u_4 = u_5 = u_6 = 0$. Thus $\pi = (5, 6)(1, 1, 1, \beta, \beta, \beta)$ and $r_2^{-1}r_2^\pi = (4, 6, 5) \in G$. This gives $\langle (1, 2, 3), (4, 5, 6) \rangle \leq G$.

Now

$$r_3^{-1}(r_3)^\pi = (2, 3)(5, 6)(\beta^{-1}, \beta^{-1}, \beta^{-1}, \beta, \beta, \beta) \in G.$$

Call this element $\hat{g}_1$. As $(1, 2, 3) \in G$, we have

$$\hat{g}_2 := \hat{g}_1^{(1,2,3)} = (1, 3)(5, 6)(\beta^{-1}, \beta^{-1}, \beta^{-1}, \beta, \beta, \beta) \in G$$

and

$$\hat{g}_3 := \hat{g}_1^{(1,3,2)} = (1, 2)(5, 6)(\beta^{-1}, \beta^{-1}, \beta^{-1}, \beta, \beta, \beta) \in G.$$

We claim that $\pi \in G^{(2)}$, from which the proof of this case immediately follows. Let $\omega = (\delta, \lambda)$ and $\omega' = (\delta', \lambda')$ be in Ω. If $\lambda, \lambda' \in \{1, 2, 3\}$, then $(\omega, \omega')^\pi = (\omega, \omega')^{g_{\omega\omega'}}$ with $g_{\omega\omega'} = 1$. If $\lambda, \lambda' \in \{4, 5, 6\}$, then $(\omega, \omega')^\pi = (\omega, \omega')^{g_{\omega\omega'}}$ with $g_{\omega\omega'} = \hat{g}_1$. Finally, suppose that only one of λ, λ' lies in $\{1, 2, 3\}$. Without loss of generality we may assume that $\lambda \in \{1, 2, 3\}$ and $\lambda' \in \{4, 5, 6\}$. Thus $\omega^\pi = (\delta, \lambda)^\pi = (\delta, \lambda)$ and $\omega'^\pi = (\delta', \lambda')^\pi = (\delta'^\beta, \lambda'^{(5,6)})$. Since $\langle c \rangle$ is transitive on Δ, there exists $x \in \langle c \rangle$ with $\delta^x = \delta^{\beta^{-1}}$. Set $g_{\omega\omega'} := \hat{g}_\lambda(x, x, x, 1, 1, 1)^{-1}$ and observe that $g_{\omega\omega'} \in G$. By construction, we have $(\omega, \omega')^\pi = (\omega, \omega')^{g_{\omega\omega'}}$.

8. Case IV: $\equiv$ has three equivalence class

Observe that the $\equiv$-equivalence classes are blocks of imprimitivity for G of size $2p$ and are union of P-orbits. In case (2) of Reduction 4.1, the group G/K has no system of imprimitivity with blocks of size 2 and hence this case cannot arise. Therefore only case (1) can happen, that is, $\sigma = 1$.

The group $G/K \cong \langle (1, 2, 3)(4, 5, 6), (1, 4)(2, 6)(3, 5) \rangle$ has three subgroups of order 2 and hence G/K has three systems of imprimitivity with blocks of size 2, namely $\{\{1, 4\}, \{2, 6\}, \{3, 5\}\}$, $\{\{1, 5\}, \{2, 4\}, \{3, 6\}\}$ and $\{\{1, 6\}, \{2, 5\}, \{3, 4\}\}$. Without loss of generality we may assume that the three $\equiv$-equivalence classes are $\Delta_1 \cup \Delta_4$, $\Delta_2 \cup \Delta_6$ and $\Delta_3 \cup \Delta_5$.

Applying Lemma 4.2 with $\rho := r_1$ and with $E \in \{\Delta_1 \cup \Delta_4, \Delta_2 \cup \Delta_6, \Delta_3 \cup \Delta_5\}$, we get

$$\hat{P} := \langle (c, 1, 1, c, 1, 1), (1, c, 1, 1, 1, c), (1, 1, c, 1, c, 1) \rangle \leq G^{(2)}.$$

Replacing π by $g^{-1}\pi$ for a suitable $g \in \hat{P}$, we may assume that $u_2 = u_3 = 0$. Furthermore, as $R_p^\pi \leq P$, we get $\ell_1 = \ell_4$, $\ell_2 = \ell_6$ and $\ell_3 = \ell_5$. It follows that $v_1 = v_4 = 0$ and $v_2 = v_6$ and $v_3 = v_5$. Write $\beta := \alpha^{v_2}$ and $\gamma := \alpha^{v_3}$. Therefore $\pi = (1, \beta, \gamma, c^{u_4}, c^{u_5}\gamma, c^{u_6}\beta)$.

We have

$$r_3^{-1}(r_3)^\pi = (c^{-u_4}, \beta^{-1}c^{-u_6}\beta, \gamma^{-1}c^{-u_5}\gamma, c^{u_4}, \gamma^{-1}c^{u_5}\gamma, \beta^{-1}c^{u_6}\beta) \in P$$

and hence $-u_4 = u_4$, $-u_5 = u_5$ and $-u_6 = u_6$. Thus $u_4 = u_5 = u_6 = 0$ and $\pi = (1, \beta, \gamma, 1, \gamma, \beta)$. Similarly, we have

$$r_2^{-1}(r_2)^\pi = (\gamma^{-1}, \beta, \beta^{-1}\gamma, \beta^{-1}, \gamma, \gamma^{-1}\beta) \in K.$$

Call this element g. As $\Delta_1 \cup \Delta_4$ is a $\equiv$-equivalence class, $\gamma^{-1} = \beta^{-1}$ and hence $\pi = (1, \beta, \beta, 1, \beta, \beta)$ and $g = (\beta^{-1}, \beta, 1, \beta^{-1}, \beta, 1)$. Applying Lemma 4.2 with $\rho := g$ and $E := \Delta_2 \cup \Delta_5$, we get $g' := (1, \beta, 1, 1, \beta, 1) \in G^{(2)}$. Thus $g'' := (g')^{r_2} = (1, 1, \beta, 1, 1, \beta) \in G^{(2)}$ and $\pi = g'g'' \in G^{(2)}$, from which the proof follows.

References

[1] L. Babai, Isomorphism problem for a class of point-symmetric structures, *Acta Math. Acad. Sci. Hungar.* **29** (1977), 329–336.

[2] W. Bosma, J. Cannon, C. Playoust, The Magma algebra system. I. The user language, *J. Symbolic Comput.* **24** (1997), 235–265.

[3] M. Conder, math review MR2335710.

[4] E. Dobson, Isomorphism problem for Cayley graphs of $\mathbb{Z}_p^3$, *Discrete Math.* **147** (1995), 87–94.

[5] C. H. Li, Z. P. Lu, P. Palfy, Further restrictions on the structure of finite CI-groups, *J. Algebr. Comb.* **26** (2007), 161–181.

[6] M. Muzychuk, On the isomorphism problem for cyclic combinatorial objects, *Discrete Math.* **197/198** (1999), 589–606.

[7] M. Muzychuk, A solution of the isomorphism problem for circulant graphs, *Proc. London Math. Soc.* **88** (2004), 1–41.

Edward Dobson, Department of Mathematics and Statistics, Mississippi State University, PO Drawer MA Mississippi State, MS 39762
E-mail address: dobson@math.msstate.edu

Joy Morris, Department of Mathematics and Computer Science, University of Lethbridge, Lethbridge, AB. T1K 3M4. Canada
E-mail address: joy@cs.uleth.ca

Pablo Spiga, Dipartimento di Matematica e Applicazioni,
University of Milano-Bicocca, Via Cozzi 55 Milano, MI 20125, Italy
E-mail address: pablo.spiga@unimib.it